

## AXIOSTAT – 100% CHITOSAN HAEMOSTATIC DRESSING ON STAB INJURY (ABDOMINAL)

### Center

Rajawadi Hospital, Mumbai-India: Dr. Sudesh Lahane

<b>Patient details</b>	<b>16 years old , Male</b>
Patient history	NA
<b>Procedure</b>	<b>Stab Injury (Abdomial)</b>
Sheath size	NA
Heparin dosage	NA
<b>Axiostat variant used</b>	<b>Axiostat Rolled Gauze, V55, L88</b>
ACT-NA	NA
<b>Average time taken to achieve haemostasis</b>	<b>8 mins</b>



Application of axiostat at the injured site

### Result

Patient outcome with Axiostat	Ease of application	Ease of removal	Adherence to wound	Conformability of dressing to wound	Patient dressing
<b>Excellent</b>					
<b>Good</b>					
<b>Fair</b>					
<b>Poor</b>					

### Discussion

- Patient was admitted to hospital with stabbed knife injury (8 inch deep) which caused lot of bleeding. Doctors tried to control bleeding for almost 2 hours by applying surgical fibrillar. 25 units of the surgical fibrillar were consumed but no hemostasis was achieved.
- Finally Axiostat was applied wnd hemostasis was achieved in 8 mins and pateint got stable